

ON GREEN AND GREEN-LAZARFELD CONJECTURES FOR SIMPLE COVERINGS OF ALGEBRAIC CURVES

E. BALLICO AND C. FONTANARI

ABSTRACT. Let X be a smooth genus g curve equipped with a simple morphism $f : X \rightarrow C$, where C is either the projective line or more generally any smooth curve whose gonality is computed by finitely many pencils. Here we apply a method developed by Aprodu to prove that if g is big enough then X satisfies both Green and Green-Lazarsfeld conjectures. We also partially address the case in which the gonality of C is computed by infinitely many pencils.

1. INTRODUCTION

Let X be a smooth complex curve of genus g . For any spanned $L \in \text{Pic}(X)$ and all integers i, j let $K_{i,j}(X, L)$ denote the Koszul cohomology groups introduced in [10]. Green's conjecture states that $K_{p,1}(X, \omega_X) = 0$ if and only if $p \geq g - \text{Cliff}(X) - 1$, where $\text{Cliff}(X)$ is the Clifford index of X , while Green-Lazarsfeld conjecture (see [11], Conjecture (3.7)) predicts that for every line bundle L on X of sufficiently large degree $K_{p,1}(X, L) = 0$ if and only if $p \geq r - \text{gon}(X) + 1$, where r is the (projective) dimension of L and $\text{gon}(X)$ is the gonality of X .

Both Green and Green-Lazarsfeld conjectures have been verified for the general curve of genus g (see [16], [17], [7], [3]) and for the general d -gonal curve of genus g (see [15] for $d \leq g/3$, [16], Corollary 1 on p. 365, for $d \geq g/3$, [7], [4]). In particular, [4] shows that both conjectures are satisfied for any smooth d -gonal curve verifying a suitable linear growth condition on the dimension of Brill-Noether varieties of pencils. Such a condition holds for the general d -gonal curve, but for special curves it turns out to be rather delicate (see [14], Statement T, and [6], Proposition 1.3).

Here we consider the case in which X is a multiple covering. Let $h : A \rightarrow B$ be a covering of degree ≥ 2 between smooth and connected projective curves. The covering h is said to be *simple* if it does not factor non-trivially, i.e. for any smooth curve D such that there are morphisms $h_1 : A \rightarrow D$ and $h_2 : D \rightarrow B$ with $h = h_2 \circ h_1$ the morphism h_2 is an isomorphism. Every covering of prime order is simple. By applying [4], Theorem 2, and [12], Theorem 1, we are going to prove the following result.

Theorem 1. *Let X be a smooth genus g curve equipped with a simple morphism $f : X \rightarrow C$ of degree $m \geq 2$, where C is a smooth curve of genus q whose gonality z is computed by finitely many g_z^1 . Assume $g \geq \max\{1 + mq + (m-1)(2mz-5), 1 + mq + (m-1)(mz-1), 4mz-9, 3mz-6\}$. Then X satisfies both Green and Green-Lazarsfeld conjectures.*

1991 *Mathematics Subject Classification.* 14H51.

Key words and phrases. Green conjecture; Green-Lazarsfeld conjecture; syzygy; covering. The authors are partially supported by MIUR and GNSAGA of INdAM (Italy).

In the special case $q = 0$, the corresponding notion of simple linear series is classical (see for instance [1]) and Green's conjecture has already been established for $m \geq 5$ in [5], Theorem 4.9, by exploiting [8] instead of [12]. By the way, for $q = 0$ our previous statement simplifies as follows.

Corollary 1. *Let X be a smooth genus g curve carrying a simple g_m^1 of degree $m \geq 3$. If $g \geq 1 + (m-1)(2m-5) \geq 4$ then X satisfies both Green and Green-Lazarsfeld conjectures.*

If instead $q > 0$ and we drop the assumption that the gonality of C is computed by finitely many pencils, we obtain with the same method the following partial result.

Proposition 1. *Let X be a smooth genus g curve equipped with a simple morphism $f : X \rightarrow C$ of degree $m \geq 2$, where C is a smooth curve of genus $q \geq 1$ and gonality $z \geq 2$. Assume $g \geq 1 + mq + (m-1)(2mz-5)$. If $m = 2$ assume also $g \geq 8z-9$. Then $K_{p,1}(X, \omega_X) = 0$ for any $p \geq g - mz + 2$ and $K_{r-mz+2,1}(C, L) = 0$ for every line bundle L on C with $h^0(C, L) = r+1$ and $\deg(L) \geq 3g$.*

Finally, if $f : X \rightarrow C$ is not simple, then $f = f_s \circ \dots \circ f_1$ with $s \geq 2$ and each f_i a simple covering. One could hope to apply Theorem 1 to each covering f_i , but the numerical restrictions on the intermediate curves make such an iterative approach effective only in very few cases.

The authors are grateful to Marian Aprodu and Claire Voisin for stimulating e-mail correspondence about Green and Green-Lazarsfeld conjectures.

2. THE PROOFS

Remark 1. Let $u : X' \rightarrow C'$ be a degree m morphism between smooth curves with X' of genus g and C' of genus q . Let $v : X' \rightarrow \mathbb{P}^1$ be a degree x morphism such that the associated morphism $(u, v) : X' \rightarrow C' \times \mathbb{P}^1$ is birational onto its image. Then $g \leq mq + (m-1)(x-1)$ (Castelnuovo-Severi inequality, see for instance [13], Corollary at p. 26). Notice that (u, v) is not birational onto its image if and only if there are a smooth curve C'' (namely, the normalization of $(u, v)(X')$) and morphisms $w : X' \rightarrow C''$, $u_1 : C'' \rightarrow C'$ and $v_1 : C'' \rightarrow \mathbb{P}^1$ such that $\deg(w) \geq 2$, $u = u_1 \circ w$ and $v = v_1 \circ w$. If u is simple, then u_1 must be an isomorphism and (u, v) is not birational onto its image if and only if there is a morphism $\eta = v_1 \circ u_1^{-1} : C' \rightarrow \mathbb{P}^1$ such that $v = \eta \circ u$. Hence in the set-up of Theorem 1 if $g \geq 1 + mq + (m-1)(mz-1)$ then X has gonality mz and for every $L \in \text{Pic}^{mz}(X)$ such that $h^0(X, L) = 2$ there is $R \in \text{Pic}^z(C)$ such that $h^0(C, R) = 2$ and $L \cong f^*(R)$.

Proof of Theorem 1. By Remark 1, X has gonality mz and $\dim(W_{mz}^1(X)) = 0$. M. Aprodu proved that X has Clifford index $mz-2$ and satisfies both Green and Green-Lazarsfeld conjectures if $\dim(W_{mz+t}^1(X)) \leq t$ for every integer t such that $0 \leq t \leq g-2mz+2$ ([4], Theorem 2). Since the function $x \rightarrow \dim(W_x^1(X))$ is strictly increasing in the interval $[\text{gon}(X), g-1]$, it is sufficient to prove $\dim(W_{g-mz+2}^1(X)) = g-2mz+2$. Assume $\dim(W_{g-mz+2}^1(X)) > g-2mz+2$, i.e. $\dim(W_{g-mz+2}^1(X)) = g-mz-j$ for some integer $j \leq mz-3$. We have $j \geq 0$ by H. Martens' Theorem ([14], see for instance [2], IV., Theorem (5.1)). Notice also that $g \geq 4j+3$ and $2j+2 \leq g-mz+2 \leq g-1-j$, hence a theorem of R. Horiuchi yields $\dim(W_{2j+2}^1(X)) = j$ ([12], Theorem 1). Let now Γ be any irreducible component of $W_{2j+2}^1(X)$ such

that $\dim(\Gamma) = j$. Since f is simple and $g - mq > (m - 1)(2mz - 5) \geq (m - 1)(2j + 1)$, by Remark 1 there are an integer $y \leq \lfloor (2j + 2)/m \rfloor$, a non-empty open subset Φ of Γ and an open subset Ψ of $W_y^1(C)$ such that every element of Φ is the pull-back of an element of Ψ plus $2j + 2 - my$ base points. Thus $j = \dim(\Gamma) = \dim(\Psi) + 2j + 2 - my \leq \dim(W_y^1(C)) + 2j + 2 - my$. We have $\Phi \neq \emptyset$, so $\Psi \neq \emptyset$ and $y \geq z$. Hence $\dim(W_y^1(C)) \leq \dim(W_z^1(C)) + 2(y - z)$ ([9], Theorem 1). Since by assumption $\dim(W_z^1(C)) = 0$, by putting everything together we get $j \leq 2(y - z) + 2j + 2 - my \leq 2j + 2 - mz$, i.e. $j \geq mz - 2$, contradiction. $\square$

The following auxiliary result provides a suitable generalization of [4], Theorem 2, by repeating almost verbatim the same proof.

Lemma 1. *Fix an integer $n \geq 1$ and let C be a smooth d -gonal curve of genus g such that $\dim G_{d+m}^1 \leq n - 1 + m$ for all m with $n - 1 \leq m \leq g - 2d + n + 1$. Then $K_{g-d+n,1}(C, \omega_C) = 0$ and $K_{r-d+n,1}(C, L) = 0$ for every line bundle L on C with $h^0(C, L) = r + 1$ and $\deg(L) \geq 3g$.*

Proof. Define integers k, ν as follows:

$$\begin{aligned} (1) \quad k &= g - d + n \\ (2) \quad \nu &= 2k - g \end{aligned}$$

and let X be the stable curve obtained from C by identifying $\nu + 1$ pairs of general points on C . In particular, let p, q be a pair of points on C identified to a node on X . If $K_{k,1}(C, \omega_C(p + q)) = 0$ then according to [7], Theorem 2.1, for every effective divisor E of degree $e \geq 1$ we have $K_{k+e,1}(C, \omega_C(p + q + E)) = 0$. Thus if L is any line bundle on C of degree $x \geq 3g$, then $h^0(C, L - \omega_C(p + q)) \geq 1$ and $K_{k+x-2g,1}(C, L) = 0$. On the other hand, by [7], Lemma 2.3 and [16], p. 367, we have $K_{k,1}(C, \omega_C) \subseteq K_{k,1}(C, \omega_C(p + q)) \subseteq K_{k,1}(X, \omega_X)$, therefore in order to prove our statement we may assume $K_{k,1}(X, \omega_X) \neq 0$ and look for a contradiction. By (2), X has genus $2k + 1$, hence by [3], Proposition 8, there exists a torsion-free sheaf F on X with $\deg(F) = k + 1$ and $h^0(X, F) \geq 2$. Let s with $0 \leq s \leq \nu + 1$ be the number of nodes at which F is not locally free. If $f : X' \rightarrow X$ is the partial normalization of X at all such nodes, then $F = f_*(L)$, where $L = f^*(F)/\text{Tors}(f^*(F))$ is a line bundle on X' with $\deg L = k + 1 - s$ and $h^0(X', L) = h^0(X, F) \geq 2$. By taking the pull-back of L on C , we obtain a g_{k+1-s}^1 not separating $\nu + 1 - s$ pairs of general points on C , hence it follows that $\dim G_{k+1-s}^1(C) \geq \nu + 1 - s$.

In order to reach a contradiction, assume first $0 \leq s \leq g - 2d + 2$ (notice that if $g = 2r - 1$ and $d = r + 1$ this case does not occur). From (1) we obtain $k + 1 - s = d - 2d + g + n + 1 - s$ with $n - 1 \leq -2d + g + n + 1 - s \leq g - 2d + n + 1$. Hence our numerical hypotheses imply that

$$\dim G_{k+1-s}^1(C) \leq g - 2d + 2n - s \leq \nu - s.$$

Assume now $s > g - 2d + 2$. We claim that also in this case

$$\dim G_{k+1-s}^1(C) = \max_r \{2(r - 1) + \dim W_{k+1-s}^r(C)\} < \nu + 1 - s.$$

Indeed, we have

$$\begin{aligned} \dim W_{k+1-s}^r(C) &\leq \dim W_{k+1-s-(r-1)}^1(C) \leq \\ &\leq \dim W_d^1(C) + 2(k + 1 - s - (r - 1) - d) \leq \\ &\leq 1 + 2(k + 1 - s - (r - 1) - d) \end{aligned}$$

where the second inequality is provided by [9], Theorem 1. Hence from (2) it follows that $\dim W_{k+1-s}^r(C) < \nu + 1 - s - 2(r - 1)$ for any r , as claimed. $\square$

Proof of Proposition 1. We argue as in the proof of Theorem 1 by applying Lemma 1 with $n = 2$ instead of [4], Theorem 2. This time we need to prove $\dim(G_{g-mz+3}^1(X)) \leq g - 2mz + 4$, so we assume by contradiction $\dim(G_{g-mz+3}^1(X)) = g - 2mz + 1 - j$ with $j \leq mz - 4$. Once again the numerical hypotheses of [12], Theorem 1, are easily checked, hence we get $j \leq \dim(W_z^1(C)) + 2j + 2 - mz$. Since in any case $\dim(W_z^1(C)) \leq 1$ by [9], Theorem 1, we obtain the desired contradiction $j \geq mz - 3$. $\square$

REFERENCES

- [1] R. D. M. Accola: On Castelnuovo's inequality for algebraic curves. I. Trans. Amer. Math. Soc. 251 (1979), 357–373.
- [2] E. Arbarello, M. Cornalba, P. A. Griffiths, and J. Harris: Geometry of algebraic curves. Vol. I. Grundlehren der Mathematischen Wissenschaften, 267. Springer-Verlag, New York, 1985.
- [3] M. Aprodu: Green-Lazarsfeld gonality conjecture for a generic curve of odd genus. Int. Math. Res. Not. 2004, no. 63, 3409–3416.
- [4] M. Aprodu: Remarks on syzygies of d -gonal curves. Math. Res. Lett. 2 (2005), 387–400.
- [5] M. Aprodu and G. Farkas: Koszul cohomology and applications to moduli, in: Aspects of vector bundles and moduli, Clay Mathematical Institute, Clay Math. Proc. Vol. 10, American Math. Soc. (2009). Pre-print arXiv:0811.3117.
- [6] M. Aprodu and G. Pacienza, The Green conjecture for exceptional curves on a $K3$ surface, Int. Math. Res. Not. IMRN 2008, no. 14, Art. ID rnn043, 25 pp.
- [7] M. Aprodu and C. Voisin: Green-Lazarsfeld's conjecture for generic curves of large gonality. C. R. Math. Acad. Sci. Paris 336 (2003), 335–339.
- [8] M. Coppens: Some sufficient conditions for the gonality of a smooth curve. J. Pure Appl. Algebra 30 (1983), 5–21.
- [9] W. Fulton, J. Harris and R. Lazarsfeld, Excess linear series on an algebraic curve, Proc. Amer. Math. Soc. 92 (1984), 320–322,
- [10] M. Green: Koszul cohomology and the geometry of projective varieties. J. Diff. Geom. 19 (1984), 125–171.
- [11] M. Green and R. Lazarsfeld: On the projective normality of complete linear series on an algebraic curve. Invent. Math. 83 (1985), 73–90.
- [12] R. Horiuchi, Gap orders of meromorphic functions on Riemann surfaces, J. Reine Angew. Math. 336 (1982), 213–220.
- [13] E. Kani, On Castelnuovo's equivalent defect, J. Reine Angew. Math. 352 (1984), 24–70.
- [14] G. Martens, On dimension theorems of the varieties of special divisors on a curve, Math. Ann. 267 (1984), 279–288.
- [15] M. Teixidor i Bigas: Green's conjecture for the generic r -gonal curve of genus $g \geq 3r - 7$. Duke Math. J. 111 (2002), 195–222.
- [16] C. Voisin: Green's generic syzygy conjecture for curves of even genus lying on a $K3$ surface. J. Eur. Math. Soc. (JEMS) 4 (2002), 363–404.
- [17] C. Voisin: Green's canonical syzygy conjecture for generic curves of odd genus. Compos. Math. 141 (2005), 1163–1190.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TRENTO, VIA SOMMARIVE 14, 38123 POVO (TN), ITALY

E-mail address: ballico@science.unitn.it, fontanar@science.unitn.it